arXiv:math/0502389v2 [math.DS] 30 May 2005

# Kolmogorov-Sinai entropy of a generalized Markov shift

IVAN WERNER
Email: ivan_werner@pochta.ru

February 1, 2008

**Abstract**

In this paper we calculate Kolmogorov-Sinai entropy $h_M(S)$ of the generalized Markov shift associated with a contractive Markov system (CMS) [11] using the coding map constructed in [12]. We show that

$$h_M(S) = -\sum_{e\in E}\int_{K_{i(e)}} p_e \log p_e d\mu$$

where $\mu$ is a unique invariant Borel probability measure of the CMS.

## 1 Introduction

In [11] we introduced a theory of *contractive Markov systems (CMS)* which provides a unifying framework in so-called "fractal" geometry. It extends the known theory of *iterated function systems (IFS)* with place dependent probabilities [1][4] in a way that it also covers *graph directed constructions* of "fractal" sets [7]. In particular, Markov chains associated with such systems also cover finite Markov chains.

By a *Markov system* we mean a family

$$\left(K_{i(e)}, w_e, p_e\right)_{e\in E}$$

(see Fig. 1) where $E$ is the set of edges of a finite directed (multi)graph $(V, E, i, t)$ ($V := \{1, ..., N\}$ is the set of vertices of the directed (multi)graph (we do not exclude the case $N = 1$), $i : E \longrightarrow V$ is a map indicating the initial vertex of each edge and $t : E \longrightarrow V$ is a map indicating the terminal vertex of each edge), $K_1, K_2, ..., K_N$ is a partition of a metric space $(K, d)$ into non-empty Borel subsets, $(w_e)_{e\in E}$ is a family of Borel measurable self-maps on the metric space such that $w_e\left(K_{i(e)}\right) \subset K_{t(e)}$ for all $e \in E$ and $(p_e)_{e\in E}$ is a family of Borel measurable probability functions on $K$ (i.e. $p_e(x) \geq 0$ for all $e \in E$ and

$\sum_{e\in E} p_e(x) = 1$ for all $x \in K$) (associated with the maps) such that each $p_e$ is zero on the complement of $K_{i(e)}$.



Fig. 1

We call a Markov system $\left(K_{i(e)}, w_e, p_e\right)_{e\in E}$ *contractive* with an *average contracting rate* $0 < a < 1$ iff it satisfies the following *condition of contractiveness on average*

$$\sum_{e\in E} p_e(x) d(w_e x, w_e y) \leq a d(x,y) \text{ for all } x, y \in K_i,\ i = 1, ..., N.$$

Markov system $\left(K_{i(e)}, w_e, p_e\right)_{e\in E}$ determines a Markov operator $U$ on the set of all bounded Borel measurable functions $\mathcal{L}^0(K)$ by

$$Uf := \sum_{e\in E} p_e f \circ w_e \text{ for all } f \in \mathcal{L}^0(K)$$

and its adjoint operator $U^*$ on the set of all Borel probability measures $P(K)$ by

$$U^*\nu(f) := \int U(f) d\nu \text{ for all } f \in \mathcal{L}^0(K) \text{ and } \nu \in P(K).$$

We say a probability measure $\mu$ is an *invariant probability measure* of the Markov system iff it is a stationary initial distribution of the associated Markov process, i.e.

$$U^*\mu = \mu.$$

A Borel probability measure $\mu$ is called *attractive* measure of the CMS if

$$U^{*n}\nu \overset{w^*}{\to} \mu \text{ for all } \nu \in P(K),$$

where $w^*$ means weak$^*$ convergence. Note that an attractive probability measure is a unique invariant probability measure of the CMS if $U$ maps continuous functions on continuous functions. We will denote the space of all bounded continuous functions by $C_B(K)$.

The main result in [11] concerning the uniqueness of the invariant measure is the following (see Lemma 1 and Theorem 2 in [11]).

**Theorem 1** *Suppose* $\left(K_{i(e)}, w_e, p_e\right)_{e\in E}$ *is an irreducible CMS such that* $K_1$,$K_2$, ...,$K_N$ *partition* $K$ *into non-empty open subsets, each* $p_e|_{K_{i(e)}}$ *is Dini-continuous and there exists* $\delta > 0$ *such that* $p_e|_{K_{i(e)}} \geq \delta$ *for all* $e \in E$*. Then:*
*(i) The CMS has a unique invariant Borel probability measure* $\mu$*, and* $\mu(K_i) > 0$ *for all* $i = 1, ..., N$*.*
*(ii) If in addition the CMS is aperiodic, then*

$$U^n f(x) \to \mu(f) \text{ for all } x \in K \text{ and } f \in C_B(K)$$

*and the convergence is uniform on bounded subsets, i.e.* $\mu$ *is an attractive probability measure of the CMS.*

A function $h : (X, d) \longrightarrow \mathbb{R}$ is called *Dini-continuous* iff for some $c > 0$

$$\int_0^c \frac{\phi(t)}{t} dt < \infty$$

where $\phi$ is *the modulus of uniform continuity* of $f$, i.e.

$$\phi(t) := \sup\{|h(x) - h(y)| : d(x, y) \leq t,\ x, y \in X\}.$$

It is easily seen that the Dini-continuity is weaker than the Hölder and stronger than the uniform continuity. There is a well known characterization of the Dini-continuity, which will be useful later.

**Lemma 1** *Let* $0 < c < 1$ *and* $b > 0$*. A function* $h$ *is Dini-continuous iff*

$$\sum_{n=0}^{\infty} \phi\left(bc^n\right) < \infty$$

*where* $\phi$ *is the modulus of uniform continuity of* $h$*.*

The proof is simple (e.g. see [11]).

Further, with the Markov system also is associated a measure preserving transformation $S : (\Sigma, \mathcal{B}(\Sigma), M) \longrightarrow (\Sigma, \mathcal{B}(\Sigma), M)$, which we call a *generalized Markov shift*, where $\Sigma := \{(..., \sigma_{-1}, \sigma_0, \sigma_1, ...) : \sigma_i \in E\ \forall i \in \mathbb{Z}\}$ is the *code space* provided with the product topology, $\mathcal{B}(\Sigma)$ denotes Borel $\sigma$-algebra on $\Sigma$ and $M$ is a *generalized Markov measure* on $\mathcal{B}(\Sigma)$ given by

$$M\left({}_m[e_1, ..., e_k]\right) := \int p_{e_1}(x) p_{e_2}(w_{e_1}x) ... p_{e_k}(w_{e_{k-1}} \circ ... \circ w_{e_1}x) d\mu(x)$$

for every thin cylinder set ${}_m[e_1, ..., e_k] := \{\sigma \in \Sigma : \ \sigma_m = e_1, ..., \sigma_{m+k-1} = e_k\}$, $m \in \mathbb{Z}$, where $\mu$ is an invariant Borel probability measure of the Markov system, and $S$ is the usual left shift map on $\Sigma$. It is easy to verify that $S$ preserves measure $M$, since $U^*\mu = \mu$ (see [12]).

For a CMS, this two pictures, Markovian and dynamical, are related by a *coding map* $F : (\Sigma, \mathcal{B}(\Sigma), M) \longrightarrow K$ which was constructed in [12]. It is defined, if $K$ is a complete metric space and $p_e|_{K_{i(e)}}$'s are Dini-continuous, by

$$F(\sigma) := \lim_{m \to -\infty} w_{\sigma_0} \circ w_{\sigma_{-1}} \circ ... \circ w_{\sigma_m} x_{i(\sigma_m)} \text{ for } M\text{-a.e. } \sigma \in \Sigma$$

where $x_i \in K_i$ for each $i = 1, ..., N$ (the coding map does not depend on the choice of $x_i$'s modulo an $M$-zero set). This coding map is the key for our calculation.

**Example 1** Let $G := (V, E, i, t)$ be a finite irreducible directed (multi)graph. Let $\Sigma_G^- := \{(..., \sigma_{-1}, \sigma_0) : \ \sigma_m \in E \text{ and } t(\sigma_m) = i(\sigma_{m-1}) \ \forall m \in \mathbb{Z} \backslash \mathbb{N}\}$ (*one-sided subshift of finite type* associated with $G$) endowed with the metric $d(\sigma, \sigma') := 2^k$ where $k$ is the smallest integer with $\sigma_i = \sigma_i'$ for all $k < i \leq 0$. Let $g$ be a positive, Dini-continuous function on $\Sigma_G$ such that

$$\sum_{y \in T^{-1}(\{x\})} g(y) = 1 \text{ for all } x \in \Sigma_G$$

where $T$ is the right shift map on $\Sigma_G^-$. Set $K_i := \left\{\sigma \in \Sigma_G^- : t(\sigma_0) = i\right\}$ for every $i \in V$ and, for every $e \in E$,

$$w_e(\sigma) := (..., \sigma_{-1}, \sigma_0, e), \ p_e(\sigma) := g(..., \sigma_{-1}, \sigma_0, e) \text{ for all } \sigma \in K_{i(e)}.$$

Obviously, maps $(w_e)_{e \in E}$ are contractions. Therefore, $\left(K_{i(e)}, w_e, p_e\right)_{e \in E}$ defines a CMS. An invariant probability measure of this CMS is called a $g$-measure. This notion was introduced by M. Keane [5]. The Kolmogorov-Sinai entropy of the generalized Markov shift, which is a natural extension of the $g$-measure $\mu$ in this case, is well known:

$$h_M(S) = \sum_{e \in E} \int g(e\sigma) \log g(e\sigma) d\mu(\sigma),$$

where $e\sigma := (..., \sigma_{-1}, \sigma_0, e)$ (see [6], [9]).

So, we are going to extend this result to a more general CMS, e.g. as in the next example.

**Example 2** Let $\mathbb{R}^2$ be normed by $\|.\|_1$. Let $K_1 := \{(x, y) \in \mathbb{R}^2 : \ y \geq 1/2\}$ and $K_2 := \{(x, y) \in \mathbb{R}^2 : \ y \leq -1/2\}$. Consider the following maps on $\mathbb{R}^2$:

$$w_1 \begin{pmatrix} x \\ y \end{pmatrix} := \begin{pmatrix} -\frac{1}{2}x - 1 \\ -\frac{3}{2}y + \frac{1}{4} \end{pmatrix}, \ w_2 \begin{pmatrix} x \\ y \end{pmatrix} := \begin{pmatrix} -\frac{3}{2}x + 1 \\ \frac{1}{4}y + \frac{3}{8} \end{pmatrix},$$

$$w_3 \begin{pmatrix} x \\ y \end{pmatrix} := \begin{pmatrix} -\frac{1}{2}|x| + 1 \\ -\frac{3}{2}y - \frac{1}{4} \end{pmatrix}, \text{ and } w_4 \begin{pmatrix} x \\ y \end{pmatrix} := \begin{pmatrix} \frac{3}{2}|x| - 1 \\ -\frac{1}{4}y + \frac{3}{8} \end{pmatrix}$$

with probability functions

$$
\begin{aligned}
p_1\begin{pmatrix} x \\ y \end{pmatrix} &:= \left(\frac{1}{15}\sin^2\|(x,y)\|_1 + \frac{53}{105}\right)1_{K_1}(x,y),\\
p_2\begin{pmatrix} x \\ y \end{pmatrix} &:= \left(\frac{1}{15}\cos^2\|(x,y)\|_1 + \frac{3}{7}\right)1_{K_1}(x,y),\\
p_3\begin{pmatrix} x \\ y \end{pmatrix} &:= \left(\frac{1}{15}\sin^2\|(x,y)\|_1 + \frac{53}{105}\right)1_{K_2}(x,y) \text{ and}\\
p_4\begin{pmatrix} x \\ y \end{pmatrix} &:= \left(\frac{1}{15}\cos^2\|(x,y)\|_1 + \frac{3}{7}\right)1_{K_2}(x,y).
\end{aligned}
$$

A simple calculation shows that $(K_{i(e)}, w_e, p_e)_{e\in\{1,2,3,4\}}$, where $i(1) = i(2) = 1$ and $i(3) = i(4) = 2$, defines a CMS with an average contracting rate 209/210. Note that none of the maps are contractive (By Theorem 2 in [11], it has a unique (attractive) invariant probability measure.)

As this paper had been known for a year as a preprint, I was informed by Wojciech Slomczynski that a similar entropy formula for some partial iterated function systems plays a central role in his recent book [8]. However, the result presented in this paper is not covered by any results from his book. Moreover, our computation is based on the existence of the coding map for some contractive Markov systems [12]. The latter is a nontrivial fact which seems to be of a fundamental nature in relation to the uniqueness of the invariant probability measure for a contractive Markov system.

## 2 Main Part

Let $\left(K_{i(e)}, w_e, p_e\right)_{e\in E}$ be a contractive Markov system with the average contracting rate $0 < a < 1$ and an invariant Borel probability measure $\mu$. We assume that: $(K, d)$ is a metric space in which sets of finite diameter are relatively compact and the family $K_1, ..., K_N$ partitions $K$ into non-empty open subsets; each probability function $p_e|_{K_{i(e)}}$ is Dini-continuous and bounded away from zero by $\delta > 0$; the set of edges $E$ is finite and the map $i : E \longrightarrow V$ is surjective. Note that the assumption on the metric space implies that it is locally compact separable and complete.

First, we prove what seems to be the main lemma for the generalized Markov shift associated with a contractive Markov system. This lemma is also used in [13]. For that we need to define some measures on the product space $K \times \Sigma$.

Denote by $\mathcal{A}$ the finite $\sigma$-algebra generated by the partition $\{_0[e] : e \in E\}$ of $\Sigma$ and define, for each integer $m \leq 1$,

$$\mathcal{A}_m := \bigvee_{i=m}^{+\infty} S^{-i}\mathcal{A},$$

which is the smallest $\sigma$-algebra containing all finite $\sigma$-algebras $\bigvee_{i=m}^{n} S^{-i}\mathcal{A}$, $n \geq m$. Let $x \in K$. For every integer $m \leq 1$, let $P_x^m$ be a probability measure on $\sigma$-algebra $\mathcal{A}_m$ given by

$$P_x^m({}_m[e_m, ..., e_n]) = p_{e_m}(x) p_{e_{m+1}}(w_{e_m}(x)) ... p_{e_n}(w_{e_{n-1}} \circ ... \circ w_{e_m}(x))$$

for all thin cylinders ${}_m[e_m, ..., e_n]$, $n \geq m$. By Lemma 1 from [12], $x \longmapsto P_x^m(Q)$ is a Borel measurable function on $K$. Therefore, we can define, for every integer $m \leq 0$,

$$\tilde{M}_m(A \times Q) := \int\limits_A P_x^m(Q)\, d\mu(x)$$

for all $A \in \mathcal{B}(K)$ and all $Q \in \mathcal{A}_m$. Then $\tilde{M}_m$ extends uniquely to a probability measure on the product $\sigma$-algebra $\mathcal{B}(K) \otimes \mathcal{A}_m$ with

$$\tilde{M}_m(\Omega) = \int P_x^m(\{\sigma \in \Sigma : (x, \sigma) \in \Omega\})\, d\mu(x)$$

for all $\Omega \in \mathcal{B}(K) \otimes \mathcal{A}_m$. Note that the set of all $\Omega \in \mathcal{B}(K) \otimes \mathcal{A}_m$ for which the integrand in the above is measurable forms a Dynkin system which contains the set all rectangles $A \times Q$, $A \in \mathcal{B}(K)$, $Q \in \mathcal{A}_m$. As the latter is $\cap$-stable and generates $\mathcal{B}(K) \otimes \mathcal{A}_m$, the integrand is measurable for all $\Omega \in \mathcal{B}(K) \otimes \mathcal{A}_m$. Further, note that $P_x^m(\{\sigma \in \Sigma : (x, \sigma) \in \Omega\}) = \int 1_\Omega(x, \sigma) dP_x^m(\sigma)$ for all $\Omega \in \mathcal{B}(K) \otimes \mathcal{A}_m$. Therefore

$$\int s d\tilde{M}_m = \int \int s(x, \sigma) dP_x^m(\sigma) d\mu(x)$$

for all $\mathcal{B}(K) \otimes \mathcal{A}_m$-simple functions $s$. Now, let $\psi$ be a $\mathcal{B}(K) \otimes \mathcal{A}_m$-measurable and $\tilde{M}_m$-integrable function on $K \times \Sigma$. Then the usual monotone approximation of positive and negative parts of $\psi$ by simple functions and the B. Levi Theorem imply that

$$\int \psi d\tilde{M}_m = \int \int \psi(x, \sigma) dP_x^m(\sigma) d\mu(x).$$

**Lemma 2** *Suppose* $C := \sum_{i=1}^{N} \int_{K_i} d(x, x_i) d\mu(x) < \infty$ *for some* $x_i \in K_i$, $i = 1, ..., N$. *Let* ${}_1[e_1, ..., e_n] \subset \Sigma$ *be a thin cylinder set . Let* $\mathcal{F} := \bigvee_{i=0}^{\infty} S^i \mathcal{A}$. *Then*

$$E_M\left(1_{{}_1[e_1,...,e_n]} | \mathcal{F}\right)(\sigma) = P_{F(\sigma)}^1\left({}_1[e_1, ..., e_n]\right) \text{ for } M\text{-a.e. } \sigma \in \Sigma,$$

*where* $E_M(.|.)$ *denotes conditional expectation with respect to measure* $M$.

*Proof.* We can obviously assume that $(e_1, ..., e_n)$ is a path of the directed graph. Set $\mathcal{F}_m := \bigvee_{i=0}^{m} S^i \mathcal{A}$ for all $m \in \mathbb{Z} \setminus \mathbb{N}$. We denote further a $(-m+1)$-tuple by $(\sigma_m, ..., \sigma_0)^*$ if $M({}_m[\sigma_m, ..., \sigma_0]) > 0$ (i.e. $(\sigma_m, ..., \sigma_0)$ is a path). Then obviously

$$E_M\left(1_{{}_1[e_1,...,e_n]} | \mathcal{F}_m\right)(\tilde{\sigma}) = \sum_{(\sigma_m,...,\sigma_0)^*} \frac{\int\limits_{{}_m[\sigma_m,...,\sigma_0]} 1_{{}_1[e_1,...,e_n]} dM}{M\left({}_m[\sigma_m, ..., \sigma_0]\right)} 1_{{}_m[\sigma_m,...,\sigma_0]}(\tilde{\sigma})$$

for $M$-a.e. $\tilde{\sigma} \in \Sigma$. Since $(\mathcal{F}_m)_{m\leq 0}$ is an increasing sequence of $\sigma$-algebras and $\mathcal{F}$ is the smallest $\sigma$-algebra containing all $\mathcal{F}_m$, it follows by Doob's Martingale Theorem (e.g. see [3] p. 199) that

$$E_M\left(1_{{}_1[e_1,...,e_n]}|\mathcal{F}_m\right) \to E_M\left(1_{{}_1[e_1,...,e_n]}|\mathcal{F}\right) \tag{1}$$

$M$-a.e..

Now, set

$$Z_m^x(\sigma) := w_{\sigma_0} \circ ... \circ w_{\sigma_m} x \text{ and } Y_m(\sigma) := w_{\sigma_0} \circ ... \circ w_{\sigma_m} x_{i(\sigma_m)}$$

for all $x \in K$, $\sigma \in \Sigma$ and $m \leq 0$. Further, define

$$X_m(\tilde{\sigma}) := \sum_{(\sigma_m,...,\sigma_0)^*} \frac{\int\limits_{K\times {}_m[\sigma_m,...,\sigma_0]} d\left(Z_m^x(\bar{\sigma}), Y_m(\bar{\sigma})\right) d\tilde{M}_m(x,\bar{\sigma})}{\tilde{M}_m\left(K \times {}_m[\sigma_m, ..., \sigma_0]\right)} 1_{{}_m[\sigma_m,...,\sigma_0]}(\tilde{\sigma})$$

for all $\tilde{\sigma} \in \Sigma$. Then

$$\begin{aligned} &\int X_m dM = \int d\left(Z_m^x(\sigma), Y_m(\sigma)\right) d\tilde{M}_m(x,\sigma) \\ = \;& \int \sum_{\sigma_m,...,\sigma_0} p_{\sigma_m}(x)...p_{\sigma_0}(w_{\sigma_{-1}}...w_{\sigma_m}x) d(w_{\sigma_0}...w_{\sigma_m}x, w_{\sigma_0}...w_{\sigma_m}x_{i(\sigma_m)}) d\mu(x) \\ \leq \;& a^{-m+1} \sum_{i=1}^{N} \int\limits_{K_i} d(x, x_i) d\mu(x) = a^{-m+1} C. \end{aligned}$$

Set

$$\Omega_m := \left\{\sigma \in \Sigma : \ X_m > a^{\frac{-m+1}{2}} C\right\}$$

for all $m \leq 0$ and $\Omega := \bigcap_{n\leq 0} \bigcup_{m\leq n} \Omega_m$. Then, by Markov inequality,

$$M\left(\Omega_m\right) \leq a^{\frac{-m+1}{2}}.$$

Hence, $M(\Omega) = 0$ and

$$X_m(\sigma) \to 0 \text{ for all } \sigma \in \Sigma \setminus \Omega.$$

Now, for $\sigma \in \Sigma$ with $M\left({}_m[\sigma_m, ..., \sigma_0]\right) > 0$,

$$\begin{aligned} &\left| \frac{\int\limits_{{}_m[\sigma_m,...,\sigma_0]} 1_{{}_1[e_1,...,e_n]} dM}{M\left({}_m[\sigma_m, ..., \sigma_0]\right)} - P^1_{F(\sigma)}\left({}_1[e_1, ..., e_n]\right) \right| \\ = \;& \left| \frac{\int p_{\sigma_m}(x)...p_{\sigma_0}(w_{\sigma_{-1}}...w_{\sigma_m}x) p_{e_1}(Z_m^x(\sigma))...p_{e_n}(w_{e_{n-1}}...w_{e_1}Z_m^x(\sigma)) d\mu(x)}{\int p_{\sigma_m}(x)...p_{\sigma_0}(w_{\sigma_{-1}}...w_{\sigma_m}x) d\mu(x)} \right. \\ & \left. - p_{e_1}(F(\sigma))...p_{e_n}(w_{e_{n-1}}...w_{e_1}F(\sigma)) \right|. \end{aligned} \tag{2}$$

Set $p(x) := p_{e_1}(x)...p_{e_n}(w_{e_{n-1}}...w_{e_1}x)$, $x \in K$. Note that the average contractiveness condition and the boundedness away from zero of the probability functions on their vertex sets imply that each map $w_e|_{K_{i(e)}}$ is Lipschitz. Since each $p_e|_{K_{i(e)}}$ is Dini-continuous, it follows that each function $p_{e_k} \circ w_{e_{k-1}} \circ ... \circ w_{e_1}|_{K_{i(e_1)}}$, $1 \leq k \leq n$, is Dini-continuous. As bounded Dini-continuous functions form an algebra, $p|_{K_{i(e_1)}}$ is also Dini-continuous. Let $\eta$ be the modulus of uniform continuity of $p|_{K_{i(e_1)}}$. By the Sublemma from [2], there exists $\psi : [0, \infty) \longrightarrow [0, \infty)$ such that $\psi(t) \geq \eta(t)$ for all $t$, $\psi(t)/t$ is non-increasing, and $\int_0^1 \psi(t)/t dt < \infty$. Set

$$\beta(u) := \frac{1}{1-a} \int\limits_0^{ua^{-1}} \frac{\psi(t)}{t} dt \text{ for } u \geq 0.$$

Then $\beta$ is continuous, concave and $\beta(0) = 0$. Moreover,

$$\beta(u) \geq \frac{1}{1-a} \int\limits_u^{ua^{-1}} \frac{\psi(t)}{t} dt \geq \frac{1}{1-a} \frac{\psi(u)}{ua^{-1}} u(a^{-1} - 1) = \psi(u) \geq \eta(u)$$

for all $u > 0$. Hence, $\eta(u) \leq \beta(u)$ for all $u \geq 0$. Therefore,

$$\begin{array}{rcl}
(2) & \leq & \left| \dfrac{\int\limits_{K \times {}_m[\sigma_m,...,\sigma_0]} p \circ Z_m^x(\bar{\sigma}) d\tilde{M}_m(x, \bar{\sigma})}{\tilde{M}_m\left(K \times {}_m[\sigma_m, ..., \sigma_0]\right)} - p \circ Y_m(\sigma) \right| + \left|p \circ Y_m(\sigma) - p \circ F(\sigma)\right| \\
& \leq & \dfrac{\int\limits_{K \times {}_m[\sigma_m,...,\sigma_0]} \left|p \circ Z_m^x(\bar{\sigma}) - p \circ Y_m(\bar{\sigma})\right| d\tilde{M}_m(x, \bar{\sigma})}{\tilde{M}_m\left(K \times {}_m[\sigma_m, ..., \sigma_0]\right)} + \left|p \circ Y_m(\sigma) - p \circ F(\sigma)\right| \\
& \leq & \dfrac{\int\limits_{K \times {}_m[\sigma_m,...,\sigma_0]} \beta\left(d\left(Z_m^x(\bar{\sigma}), Y_m(\bar{\sigma})\right)\right) d\tilde{M}_m(x, \bar{\sigma})}{\tilde{M}_m\left(K \times {}_m[\sigma_m, ..., \sigma_0]\right)} + \left|p \circ Y_m(\sigma) - p \circ F(\sigma)\right| \\
& \leq & \beta\left(\dfrac{\int\limits_{K \times {}_m[\sigma_m,...,\sigma_0]} d\left(Z_m^x(\bar{\sigma}), Y_m(\bar{\sigma})\right) d\tilde{M}_m(x, \bar{\sigma})}{\tilde{M}_m\left(K \times {}_m[\sigma_m, ..., \sigma_0]\right)}\right) + \left|p \circ Y_m(\sigma) - p \circ F(\sigma)\right| \\
& = & \beta \circ X_m(\sigma) + \left|p \circ Y_m(\sigma) - p \circ F(\sigma)\right|.
\end{array}$$

Hence

$$\left|E_M\left(1_{{}_1[e_1,...,e_n]}|\mathcal{F}_m\right)(\sigma) - P^1_{F(\sigma)}\left({}_1[e_1, ..., e_n]\right)\right| \leq \beta \circ X_m(\sigma) + \left|p \circ Y_m(\sigma) - p \circ F(\sigma)\right|$$

for $M$-a.e. $\sigma \in \Sigma$. By Corollary 1 from [12] and the continuity of $p$ on $K$, the second term also converges to zero $M$-a.e.. Thus

$$\left|E_M\left(1_{{}_1[e_1,...,e_n]}|\mathcal{F}_m\right)(\sigma) - P^1_{F(\sigma)}\left({}_1[e_1, ..., e_n]\right)\right| \to 0 \text{ as } m \to -\infty$$

for $M$-a.e. $\sigma \in \Sigma$. With (1), this implies that

$$E_M \left( 1_{{}_1[e_1,...,e_n]} | \mathcal{F} \right)(\sigma) = P^1_{F(\sigma)} \left( {}_1[e_1, ..., e_n] \right)$$

for $M$-a.e. $\sigma \in \Sigma$. $\square$

**Proposition 1** *If invariant probability measure $\mu$ is unique, then*

$$F(M) = \mu.$$

*Proof.*Let $U^*$ be the adjoint of the Markov operator associated with the CMS. It is sufficient to show that $U^*F(M) = F(M)$, since $\mu$ is the unique invariant probability measure. Let $f \in C_B(K)$. Then

$$U^*F(M)(f) \quad = \quad \sum_{e \in E} \int p_e f \circ w_e dF(M) = \sum_{e \in E} \int p_e \circ F f \circ w_e \circ F dM.$$

Let $e \in E$. By Theorem 1 (iv) from [11], $\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty$. Therefore, by Lemma 2,

$$E_M \left( 1_{{}_1[e]} \middle| \mathcal{F} \right)(\sigma) = P^1_{F(\sigma)} \left( {}_1[e] \right) = p_e \circ F(\sigma)$$

for $M$-a.e. $\sigma \in \Sigma$. Since $f \circ w_e \circ F$ is bounded and $\mathcal{F}$-measurable, it follows by a well known property of the conditional expectation that

$$E_M \left( 1_{{}_1[e]} f \circ w_e \circ F \middle| \mathcal{F} \right)(\sigma) = p_e \circ F(\sigma) f \circ w_e \circ F(\sigma)$$

for $M$-a.e. $\sigma \in \Sigma$. Hence, by the shift invariance of $M$,

$$\begin{aligned} U^*F(M)(f) &= \sum_{e \in E} \int 1_{{}_1[e]}(\sigma) f \circ w_e \circ F(\sigma) dM(\sigma) \\ &= \sum_{e \in E} \int 1_{{}_1[e]} \circ S^{-1}(\sigma) f \circ w_e \circ F \circ S^{-1}(\sigma) dM(\sigma) \\ &= \sum_{e \in E} \int 1_{{}_0[e]}(\sigma) f \circ F(\sigma) dM(\sigma) \\ &= F(M)(f). \end{aligned}$$

$\square$

**Theorem 2** *Let $h_M(S)$ be Kolmogorov-Sinai entropy of the generalized Markov shift associated with the contractive Markov system.*
*(i) If $\sum_{i=1}^N \int_{K_i} d(x, x_i) d\mu(x) < \infty$ for some $x_i \in K_i$, $i = 1, ..., N$, then*

$$h_M(S) = -\sum_{e \in E} \int_{K_{i(e)}} p_e \log p_e dF(M).$$

$(ii)$ *If invariant probability measure* $\mu$ *is unique, then*

$$h_M(S) = -\sum_{e\in E}\int_{K_{i(e)}} p_e \log p_e d\mu.$$

*Proof.* It is well known that $h_M(S) = h_M(S^{-1})$ (e.g. Theorem 4.13 in [10]) and, by the Kolmogorov-Sinai Theorem (e.g. Theorem 4.17 in [10]), $h_M(S^{-1}) = h_M(S^{-1}, \mathcal{A})$. Further, using the notion of conditional entropy (e.g. Theorem 4.3 (ix) and Theorem 4.14 in [10]),

$$h_M(S^{-1}, \mathcal{A}) = \mathcal{H}\left(S^{-1}\mathcal{A} \middle/ \bigvee_{i=1}^{\infty} S^{i-1}\mathcal{A}\right).$$

Set $\mathcal{F} := \bigvee_{i=0}^{\infty} S^i\mathcal{A}$. Hence

$$h_M(S) = -\sum_{e\in E}\int E\left(1_{{}_1[e]}\middle|\mathcal{F}\right)\log E\left(1_{{}_1[e]}\middle|\mathcal{F}\right)dM.$$

By the assumption in $(i)$, Lemma 2 implies that

$$E\left(1_{{}_1[e]}\middle|\mathcal{F}\right) = p_e \circ F \ M\text{-a.e.}$$

for each $e \in E$. Hence, with $0\log 0 = 0$, we have

$$h_M(S) = -\sum_{e\in E}\int p_e \circ F \log\left(p_e \circ F\right) dM = -\sum_{e\in E}\int_{K_{i(e)}} p_e \log p_e dF(M).$$

By the assumption in $(ii)$, Theorem 1(iv) in [11] and Proposition 1 imply that

$$h_M(S) = -\sum_{e\in E}\int_{K_{i(e)}} p_e \log p_e d\mu.$$

□

## Acknowledgements

This work was done as a part of my PhD thesis at the University of St Andrews. I would like to thank: EPSRC and School of Mathematics and Statistics of University of St Andrews for providing me with a scholarship and excellent working conditions in St Andrews, Professor K. J. Falconer for valuable comments on the first draft of the paper, my supervisor Lars Olsen for his interest in my work, his support and many fruitful discussions.

# References

[1] M. F. Barnsley, S.G.Demko, J.H.Elton and J.S.Geronimo, Invariant measure for Markov processes arising from iterated function systems with place-dependent probabilities, *Ann. Inst. Henri Poincaré* **24** $n^{\circ}3$ (1988) 367-394.

[2] M. F. Barnsley, S.G.Demko, J.H.Elton and J.S.Geronimo, Erratum; Invariant measure for Markov processes arising from iterated function systems with place-dependent probabilities, *Ann. Inst. Henri Poincaré***25** $n^{\circ}4$ (1989) 589-590.

[3] J. L. Doob, *Measure Theory*, Spriger-Verlag (1993).

[4] J. H. Elton, An ergodic theorem for iterated maps, *Ergod. Th. & Dynam. Sys.* **7** (1987) 481-488.

[5] M. Keane, Strongly Mixing $g$-Measures, *Inventiones math.* **16** (1972) 309-324.

[6] F. Ledrappier, Principe variationnel et systèmes dynamiques symboliques, *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **30** (1974), 185-202.

[7] R. D. Mauldin and S. C. Williams, Hausdorff dimension in graph directed constructions, *Tran. AMS* **309** (1988) 811-829.

[8] W. Slomczynski, *Dynamical entropy, Markov operators, and itereated function systems*, Rozprawy Habilitacyjne Uniwersytetu Jagiellońskiego Nr 362, Wydawnictwo Uniwersytetu Jagiellońskiego (2003).

[9] P. Walters, Ruelle's Operator Theorem and $g$-measures, *Tran. AMS* **214** (1975) 375-387.

[10] P. Walters, *An introduction to Ergodic Theory*, Springer-Verlag, (1982).

[11] I. Werner, Contractive Markov systems, *J. London Math. Soc.*. **71** (2005) 236-258.

[12] I. Werner, Coding map for a contractive Markov system, *Math. Proc. Camb. Phil. Soc.* to appear **140** (2006).

[13] I. Werner, Ergodic theorem for a contractive Markov system, *Nonlinearity* **17** (2004) 2303-2313.